

Clonal effect on rooting and acclimation rates for *in-vitro* micropropagation in hybrid walnut (*Juglans x intermedia* Mj 209): preliminary observations

Jesus Fernández-Moya^{1*}, Ricardo Julian Licea-Moreno¹, Ignacio Urbán-Martínez¹, Ruth M. Castro-Fernández¹, Cristina Ramírez-López-Ramallal¹

Received 3/09/2019 - Accepted 4/01/2020 - Published online 07/02/2020

Abstract - The success of walnut (*Juglans* sp.) planted forests for timber production have been very variable and genetic material is considered as one of the main drivers (together with site selection and forest management) for the success or failure of the plantations, as the performance of the trees from seed material is very variable. Considering the relevance of this genetic material, several clones have been selected and research have been conducted in order to improve micropropagation procedures. The objective of the present study is to analyze the effects of different clones in the rooting and acclimation rates for *in-vitro* micropropagation in hybrid walnut (*Juglans x intermedia* Mj 209). The results show a significant effect of clones on the rooting and the total micropropagation efficiency rates, but not on the acclimation rate. The efficiency rate of D-117 (65%) is considered statistically higher than the one for D-15 (38%), caused by a higher rooting rate of D-117 (73%) compared with D-15 (55%), because acclimation rate (57%) did not show any clone effect. Considering these differences in the micropropagation success, it might be considered (together with other factors) for clone selection to increase the general performance of the plant production units in large-scale propagation.

Keywords - clone selection, hybrid walnut, micropropagation, vitroplants, rooting, acclimation.

Introduction

Walnut trees are species of the genus *Juglans* spp. L., traditionally characterized by their highly-valued nuts and timber. Considering the high timber value and the shortage of the species, many walnut forest plantations oriented for timber production have been established during the last decades (Mohani et al. 2009). These plantations have been established with the common species of Persian or European walnut (*J. regia* L.) and Black or American walnut (*J. nigra* L., *J. major* (Torr.) A. Heller, *J. hindsii* (Jeps.) Jeps. ex R.E. Sm.) but also with several hybrids which have been specifically developed for timber production, e.g.: Mj-209xRa y Ng-23xRa (Aletà 2004, Victory et al. 2006, Mohani et al. 2009, Clark and Hemery 2010, Coello et al. 2013).

The success of this kind of planted forests have been very variable and the genetic material is considered as one of the main drivers (together with site selection and forest management) for the success or failure of the plantations (e.g. Aletà et al. 2003, Aletà 2004, Urbán-Martínez et al. 2013, Licea-Moreno 2016). To this respect, Aletà and Vilanova (2006) pointed out the high variability in the performance of the trees from seed material, despite the trees are

genetically close between them. Considering the relevance of this genetic material selection for walnut (*Juglans* spp.) planted forests oriented for timber production, there have been two big research groups working in the clone selection in Spain: 1) from the public center IRTA (e.g. Aletà et al. 2003, Aletà 2004) and 2) from the private company Bosques Naturales SA (e.g. Urbán-Martínez et al. 2013, Licea-Moreno 2016). To this respect, the Spanish Register for Forest Reproductive Material includes 29 clones (21 for *J. nigra*, 8 for *J. regia* and 10 for *J. x intermedia*), 9 parents of family (5 for *J. regia* and 4 for *J. x intermedia*) and one seed orchard of *J. regia* (MAPA 2016).

Urbán-Martínez et al. (2013) published a detailed characterization of the selection process of the Bosques Naturales SA clones. At the moment, the *in-vitro* micropropagation plant production is focused in 4 clones (D-15, D-117, D-53 and D-M) selected by their vigour, trunk straightness (based on the classification of MacDonald et al. [2001]) and wood quality. The clone D-15 (commercial name NAT-7-BN) is the one most widely used by the company because it was one of the firsts to be introduced *in vitro*. The original D-15 plus tree is in Cáceres and it had, with 12 years old, 13.5 m height,

¹ Bosques Naturales SA, Alcobendas, Madrid – Spain

*Corresponding author: jesusfernandez@bosquesnaturales.com

20.6 cm DBH, 3 m of clear boles and a straightness score of 2/7. Fernández-Moya et al. (in press) analyze the field performance of the planted forests established with the clone NAT-7-BN in their forests at A Coruña, Toledo, Cuenca and Girona. The original D-117 plus tree is in Girona and it had, with 12 years old, 17.3 m height, 29.5 cm DBH, 5 m of clear boles and a straightness score of 7/7. The original D-53 plus tree is in Cáceres and it had, with 13 years old, 18.1 m height, 24.4 cm DBH, 5 m of clear boles and a straightness score of 4/7. The original D-M plus tree is in Cáceres and it had, with 13 years old, 15.2 m height, 22.6 cm DBH, 4 m of clear boles and a straightness score of 7/7.

Despite the interest on walnut clones, traditional vegetative methods are not suitable for massive reproduction of walnut elite genotypes; becoming tissue culture technologies the most important alternative for cloning, but the high recalcitrance of walnut species hinder their commercial micropropagation (Licea-Moreno 2016). Since the first *in-vitro* procedures were developed for walnut (Driver and Kuniyuki 1984, McGranahan et al. 1987, McGranahan and Leslie 1988, Cornu and Jay-Allemand 1989), a number of modifications have been tested to increase success and extend the methodology to different species and hybrids (e.g., Dolcet-Sanjuan et al. 2004, Leal et al. 2007, Bosela and Michler 2008, Leslie et al. 2009, Vahdati et al. 2009, Toosi and Dilmagani 2010, Licea-Moreno et al. 2012 and 2015, Licea-Moreno 2016). In addition to the problem of introducing field-selected material to *in-vitro* culture, better procedures are needed regarding microshoot multiplication, microshoot rooting and plantlet acclimatization. Poor rooting has been reported as arguably the main factor limiting the establishment of clonal plantations (Woeste and Michler 2011).

As a consequence of these difficulties for the production of clonal plants in the nurseries, there is a shortage of clonal plant material in the market. To this respect, Bosques Naturales SA is one of the key players regarding clonal walnut planted forests in Europe. Since 1998, an experimental micropropagation protocol was developed (Licea-Moreno et al. 2012 and 2015, Licea-Moreno 2016) and the company has been producing around 2,000 vitroplants per year (high quality clonal plantlets) in their research facilities. The objective of the present study is to analyze the effects of the different clones in the rooting and acclimation rates for *in-vitro* micropropagation in hybrid walnut (*Juglans x intermedia* Mj 209) using the available data of the annual production of the company Bosques Naturales SA.

Material and Methods

Experimental design: clones used and data collection

The present study is based on the analysis of the work during the 2017 plant production campaign (from 22/December/2016 to 6/November/2017) in the Bosques Naturales SA facilities in Madrid (*in-vitro* laboratory) and Galicia (nursery for acclimation). Hence, there is not a fixed and pure experimental design for this work and the available data of the annual production of the company is used for the analysis.

Within the pool of clones registered by Bosques Naturales SA (all of them of the hybrid walnut -*Juglans x intermedia* Mj 209), 4 clones were selected for the analysis: D-117, D-15, D-53 and D-M. All the selected clones are registered at the Spanish Register for Forest Reproductive Material (MAPA 2016). These clones come from a selection of walnut elite genotypes and introduced *in-vitro* in 2008/2009. More details of the selection and *in-vitro* introduction procedures have been previously published (Licea-Moreno et al. 2012, Urbán-Martínez et al. 2013, Licea-Moreno et al. 2015, Licea-Moreno 2016).

The data collection is derived from the daily/weekly control registered in the facilities. To this respect, the micropropagation is performed by batches of plants. There were 40 batches analysed: 16 from D-117 (2,356 explants), 14 from D-15 (1,455 explants), 5 from D-53 (345 explants) and 5 from D-M (299 explants).

In-vitro micropropagation protocol

The specific protocol developed by Bosques Naturales for walnut (*Juglans* sp.) *in-vitro* micropropagation has been published in detail in Licea-Moreno et al. (2012 and 2015) Licea-Moreno (2016). This protocol is divided into several stages: multiplication, elongation, pre-induction, rooting and acclimation. The present study is focused on the final stages of the protocol, which are described with more detail as follows.

Pre-induction stage.

This consists in 5 days of total darkness immersed in a medium (6 ml per explant) containing: 30 g L⁻¹ of saccharose, stock solutions: 50% of macronutrients and 100% of micronutrients and vitamins from prepared DKW formula, 119 mg L⁻¹ of FeEDHA and considerable higher dose of IBA, 10 mg L⁻¹.

Table 1 - ANOVA results of the effect of clones on the rooting rate (%), acclimation rate (%) and efficiency (%) of hybrid walnut (*Juglans x intermedia* Mj209) in vitro micropropagation.

Variable	F - value	P - value
Rooting rate	9.016	0.000138
Acclimation rate	1.05	0.383
Efficiency	2.567	0.0701

Rooting stage.

Microshoots from radical pre-induction stay for 2 to 3 weeks, depending on genotype, in a preparation of vermiculite (instead of agar), intermediate granulometry 0.5-3 mm, soaked with medium made off with 16.5 g L⁻¹ of glucose, 119 g L⁻¹ FeEDDHA and stock solutions: 50% of macronutrients and 100% of micronutrients and vitamins from prepared DKW formula. Cristal vessel are used with 10 explants per vessel with 80 g of vermiculite and 100 mL of medium, although this relation must be adjusted depending on each type of vessel.

Acclimation stage.

Finally, acclimation phase takes place for those explants that have developed roots in previous stage preferably. It will last in this stage for 4 weeks, first 2 weeks guaranteeing in a greenhouse a minimum of 75% of relative humidity (rh) and a lighting of 150 μmol m⁻² s⁻¹, the following 3rd and 4th week 70% of rh and 150 μmol m⁻² s⁻¹. To transplant the explants to pots for this stage, it is preferable to use thin black and blonde peat (50%-50%) mixed with one part of vermiculite.

Statistical analysis

Rooting rate (%), acclimation rate (%) and efficiency (%) were analyzed, defined as follow

Rooting rate (%) = rooted microshoots / initial explants

Acclimation rate (%) = acclimated vitroplants / rooted microshoots

Efficiency (%) = acclimated vitroplants / initial explants

An ANOVA was performed to test the effect of clones on each of these variables and a Tukey's HSD test was also done when this effect was identified as significant. General Linear Models were used in all the cases as the diagnostic checking performed showed that the data respect the model assumptions. A significance level of 0.1 is considered if the contrary is not stated. All the statistical analyses were performed using R (R Core Team 2019).

Results and Discussion

Rooting stage

The results show a significative effect of clones on the rooting rate in the micropropagation process for hybrid walnut (Fig. 1, Tab. 1). The model was considered as adequate after a normality analysis of the residuals (Shapiro test p-value=0.5764) and their graphical diagnosis (Fig. 2). The rooting rate of D-117 [73% ± 7 (C.I. 90%)] is considered statistically similar than the one for D-53 [93% ± 5 (C.I. 90%)] and D-M [88% ± 6 (C.I. 90%)], while the rate for D-15 [55% ± 9 (C.I. 90%)] is significantly lower (Fig. 1)




Figure 1 - Effect of clones on the rooting rate (%), acclimation rate (%) and efficiency (%) of hybrid walnut (*Juglans x intermedia* Mj209) in vitro micropropagation.

The effect of different clones on the rooting rate is a common pattern which has been detected either for walnut species (*Juglans* spp.) (Chenevard et al. 1995, Dolcet-Sanjuan et al. 1996, Scaltsoyiannes et al. 1997, Vahdati et al. 2004, Sharifian et al. 2009, Payghanzadeh and Kazemitabar 2011) and other hardwoods (Bennett and McComb 1982, San José et al. 1988, Juncker and Favre 1989, Yu et al. 2001), while Tetsumura et al. (2002) did not find a significant effect on the rooting of *J. regia* explants. To this respect, the rooting rates reported for the different clones (Fig. 1) are within the range reported




Figure 2 - Graphical evaluation of the residuals of the statistical model analyzing the effect of clones on the rooting rate (%) of hybrid walnut (*Juglans x intermedia* Mj209) in vitro micropropagation.

for the *Juglans* spp. species by other authors (Che-nevard et al. 1995, Dolcet-Sanjuan et al. 1996, Scaltsoyiannes et al. 1997, Tetsumura et al. 2002, Vahdati et al. 2004, Sharifian et al. 2009)

Acclimation stage

Notwithstanding the clone effect on the rooting rate, the results do not show a significant effect of clones on the acclimation rates in the micropropagation process for hybrid walnuts (Tab. 1, Fig. 1). Hence, regardless the clone considered, the acclimation rate is $57\% \pm 7$ (C.I. 90%). This poor acclimation results have been previously considered as a key issue for the success in the operational micropropagation of walnut, as the acclimatization of micropropagated walnut is reported to be a difficult procedure because of rapid desiccation of plantlets or their susceptibility to diseases due to high humidity and difficult rooting, for a review see Payghamzadeh and Kazemitabar (2011). As oppose as the results from this study, the clonal effect on the acclimation rates of walnut vitroplants has been previously reported by other authors (e.g. Dolcet-Sanjuan et al. 1996, Frossard et al. 1997). Indeed, Frossard et al. (1997) report how *J. regia* clones are more easily adapted to soil conditions than *J nigra* x *J regia* hybrid clones. Improving this acclimation stage is considered as a priority for the commercial walnut micropropagation and there have been many authors reporting several measures to improve the acclimation rates (e.g. Jay-Allemand et al. 1992, Heloir et al. 1994, Voyiatzis and Mc Granahan 1994, Dolcet-Sanjuan et al. 1996, Frossard et al. 1997). To this respect, the acclimation rates of the 2018 cam-

paign are higher than 65% on average, even though the data is not shown in this work due to big differences in the experimental conditions.

Efficiency

Based on the rooting and acclimation rates, the results show a significant effect of clones on the micropropagation efficiency for hybrid walnuts (Tab. 1, Fig. 1). The model was considered as adequate after a normality analysis of the residuals (Shapiro test p-value=0.3744) and their graphical diagnosis (Fig. 3). The efficiency rate of D-117 [$65\% \pm 15$ (C.I. 90%)] is considered statistically higher than the one for D-15 [$38\% \pm 10$ (C.I. 90%)], while the high variability of the clones D-53 [$50\% \pm 17$ (C.I. 90%)] and D-M [$51\% \pm 13$ (C.I. 90%)] cause that they cannot be considered as different from any of them (Fig. 1). Considering these differences in the micropropagation efficiency, clones might be selected in order to improve the general performance of the plant production units in large-scale propagation, as suggested by other authors (e.g. Scaltsoyiannes et al. 1997). However, multiplication rates are not considered in the present study and it is a key issue in order to be considered for this selection in addition to other key issues such as field performance (e.g. Aletà et al. 2003, Urbán-Martínez et al. 2013). To this respect, D-15 and D-117 show a better multiplication rate (data not shown) which partially explains why they are the two clones more used in this study (and in the company's commercial production).




Figure 3 - Graphical evaluation of the residuals of the statistical model analyzing the effect of clones on the efficiency rate (%) of hybrid walnut (*Juglans x intermedia* Mj209) *in vitro* micropropagation.

Conclusion

The results show a significant effect of clones on the rooting and the total micropropagation efficiency rates of the hybrid walnut (*Juglans x intermedia* Mj 209), but not on the acclimation rate. The efficiency rate of D-117 [65%±15 (C.I. 90%)] is considered statistically higher than the one for D-15 [38%±10 (C.I. 90%)], caused by a higher rooting rate of D-117 [73%±7 (C.I. 90%)] compared with D-15 [55%±9 (C.I. 90%)], because acclimation rate [57%±7 (C.I. 90%)] did not show any clone effect. Considering these differences in the micropropagation success, it might be considered (together with other factors) for clone selection to increase the general performance of the plant production units in large-scale propagation.

Acknowledgements

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 728086: Innovation Action "Woodnat" (Second generation of planted hardwood forests in the EU).

References

- Aletà N., Ninot A., Voltas J. 2003 - *Caracterización del comportamiento agroforestal de doce genotipos de nogal (Juglans spp.) en dos localidades de Cataluña*. Investigación Agraria: Sistemas y Recursos Forestales 12 (1): 39-50.
- Aletà N. 2004 - *Current Research in Spain on Walnut for Wood Production*. In: Proceedings 6th Walnut Council research symposium Michler et al. (Eds.) "Black walnut in a new century"; Lafayette; U.S. Department of Agriculture, Forest Service, North Central Research Station. 188 p.
- Aletà N., Vilanova A. 2006 - *El nogal híbrido*. Navarra Forestal 13:18-21.
- Bennett I.J., McComb J.A. 1982 - *Propagation of jarrah (Eucalyptus marginata Donn. ex Sm.) by organ and Tissue culture*. Australian Forest Research 12: 121-127.
- Bosela M.J., Michler C.H. 2008 - *Media effects on black walnut (Juglans nigra L.) shoot growth in vitro: evaluation of multiple nutrient formulations and cytokinin types*. In vitro Cellular & Development Biology - Plant 44: 316-329.
- Chenevard D., Jay-Allemand C., Gendraud M., Frossard J.S. 1995 - *The effect of sucrose on the development of hybrid walnut microcuttings (Juglans nigra x Juglans regia). Consequences on their survival during acclimatization*. Annales des Sciences Forestières 52 (2): 147-156.
- Clark J., Hemery G. 2010 - *Walnut hybrids in the UK: fast growing quality hardwoods*. Quarterly Journal of Forestry 104: 43-46.
- Coello J., Becquey J., Gonin P., Ortisset J.P., Desombre V., Baiges T., Piqué M. 2013 - *Guía de ecología y silvicultura de especies productoras de madera de calidad*. Generalidad de Cataluña - Centro de la Propiedad Forestal.
- Cornu D., Jay-Allemand C. 1989 - *Micropropagation of hybrid walnut trees (Juglans nigra x Juglans regia) through culture and multiplication of embryos*. Annals of Forest Science 46S: 113-116.
- Driver J.A., Kuniyuki A.H. 1984. *In vitro propagation of paradox walnut rootstock*. HortScience 19 (4): 507-509.
- Dolcet-Sanjuan R., Claveria E., Camprubí A., Estaún V., Calvet C. 1996 - *Micropropagation of walnut trees (Juglans regia L.) and response to arbuscular mycorrhizal inoculation*. Agronomie 16: 639-645.
- Dolcet-Sanjuan R., Claveria E., Gruselle R., Meier-Dinkel A., Jay-Allemand C., Gaspar T. 2004 - *Practical factors controlling in-vitro adventitious root formation from walnut shoots microcuttings*. Journal of the American Society for Horticultural Science 129:198-203.

- Fernández-Moya J., Licea-Moreno R., Santacruz D., Urbán-Martínez I. (in press) - *NAT 7 BN: bosques plantados clonales de nogal híbrido (Juglans x intermedia MJ 209) para la producción sostenible de maderas nobles*. Accepted for publication on the journal Cuadernos de la Sociedad Española de Ciencias Forestales.
- Frossard J.S., Dolcet-Sanjuan R., Chenevard D., Jay-Allemand C. 1997 - *Physiological aspects of microcuttings during acclimatization*. In: "European Development of Walnut Trees for Wood and Fruit Production as an Alternative and Extensive System to Agricultural Crops". C Jay-Allemand, ed., European Commission Directorate-General VI, Luxemburg.
- Heloir M.C., Kevers C., Hausman J.F., Deltour R., Gaspar T. 1994 - *Histological origin of roots of micropropagated walnut (Juglans regia L.) shoots*. Archives Internationales de Physiologie, de Biochimie et de Biophysique 102.
- Jay-Allemand C., Capelli P., Cornu D. 1992 - *Root development of in vitro hybrid walnut microcuttings in a vermiculite-containing gelrite medium*. Scientiae Horticulturae 51: 335-342.
- Juncker B., Favre J.M. 1989 - *Clonal effects in propagating oak trees via in vitro culture*. Plant Cell Tissue and Organ Culture 19 (3): 267-276.
- Leal D.R., Sánchez-Olate M., Avilés F., Materan M.E., Uribe M., Hasbún R., Rodríguez R. 2007 - *Micropropagation of Juglans regia L.* In: "Protocols for micropropagation of woody trees and fruits". Jain S.M., Häggman H. (eds) Springer, Heidelberg: 381-390.
- Leslie C.A., Hackett W.P., McGranahan G.H. 2009 - *Improved rooting methods for walnut (Juglans) microshoots*. Acta Horticulturae 861:365-372.
- Licea-Moreno R.J. 2016 - *Biotecnología forestal aplicada a la producción de madera de nogal*. Universidad Politécnica de Madrid. PhD Thesis.
- Licea-Moreno R.J., Morales A.V., Daquinta M., Gómez L. 2012 - *Towards scaling-up the micropropagation of Juglans major (Torrey) Heller var. 209 x J. regia L., a hybrid walnut of commercial interest*. In: Proceedings of the IUFRO Working party 2.09.02 conference on "Integrating vegetative propagation, biotechnologies and genetic improvement for tree production and sustainable forest anagement" Park, Y.S. and Bonga, J.M. (eds). Brno Czech Republic, June 25-28. [Online] Available: <http://www.iufro20902.org/> : 80-91.
- Licea-Moreno R.J., Contreras A., Morales A.V., Urban I., Daquinta M., Gomez L. 2015 - *Improved walnut mass micropropagation through the combined use of phloroglucinol and FeEDDHA*. Plant Cell Tissue and Organ Culture 123 (1):143-154.
- MAPA 2016 - *Informe anual: Situación del Registro y Catálogo Nacional de Materiales de Base*. Ministerio de Agricultura, Pesca y Alimentación, Gobierno de España. [Online] Available: www.mapa.gob.es/es/desarrollo-rural/temas/politica-forestal/informeanual2016_situaciondelregistrocnmb_tcm30-139340.pdf
- Macdonald E.S., Mochan S., Connolly T. 2001 - *Protocol for Stem Straightness Assessment in Sitka Spruce*. Information Note. Forestry Commission 231 Corstorphine Road. Edinburgh. EH12 7AT.
- McGranahan G., Leslie C.A. 1988 - *In-vitro propagation of mature Persian walnut cultivars*. Horticultural Science 23: 220.
- McGranahan G.H., Driver J.A., Tulecke W. 1987 - *Tissue culture of Juglans*. In: "Cell and tissue culture in forestry" Vol. 3 Bonga JM, Durzan DJ (eds). Martinus Nijhoff, Boston: 261-271.
- Mohani C., Pelleri F., Hemery G.E. 2009 - *The modern silviculture of Juglans regia L.: A literature review*. Die Bodenkultur 60 (3): 21-34.
- Payghamzadeh K., Kazemitabar S.K. 2011 - *In vitro propagation of walnut - A review*. African Journal of Biotechnology 10 (3): 290-311.
- R Core Team 2019 - *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. Vienna, Austria. [Online] Available: <http://www.R-project.org>.
- San José M.C., Ballester A., Vieitez A.M. 1988 - *Factors affecting in vitro propagation of Quercus robur L.* Tree Physiology 4 (3): 281-290.
- Scaltsioyianes A., Tsoulpha P., Panetsos K.P., Moulalis D. 1997 - *Effect of Genotype on Micropropagation of Walnut Trees (Juglans regia L.)*. Silvae Genetica 46 (6): 326-332.
- Sharifian S., Vahdati K., Mirmasoumi M., Ghaem Maghami S.A. 2009 - *Assessment of Phloroglucinol Effect on Rooting of Tissue Cultured Persian Walnut*. In: Proceedings III IS on Acclim. and Establ. of Micropropagated Plants. Acta Horticulturae 812. Ed. A. Romano.
- Tetsumura T., Tsukuda K., Kawase K. 2002 - *Micropropagation of Shinano Walnut (Juglans regia L.)*. Journal of Japanese Society for Horticultural Science 71 (5): 661-663.
- Toosi S., Dilmagani K. 2010 - *Proliferation of Juglans regia L. by in vitro embryo culture*. Journal of Cell Biology and Genetics 1: 12-19.
- Urbán-Martínez I., Licea-Moreno R.J., González C., Villar de Arcos F., Santaclara O., Merlo-Sánchez E. 2013 - *Characterization of walnut hybrid plus trees for good quality wood plantations*. In: Proceedings of International Scientific Conference on Hardwood (ISCHP2013), Florencia
- Vahdati K., Leslie C., Zamani Z., McGranahan G. 2004 - *Rooting and acclimatization of in vitro-grown shoots from mature trees of three persian walnut cultivars*. Horticultural Science 39: 324-327.
- Vahdati K., Razaee R., Mirmasoomi M. 2009 - *Micropropagation of some dwarf and early mature walnut genotypes*. Biotechnology 8:171-175.
- Victory E.R., Glaubitz J.C., Rhodes O.E., Woeste K.E. 2006 - *Genetic homogeneity in Juglans nigra L. (Juglandaceae) at nuclear microsatellites*. American Journal of Botany 93:118-126.
- Voyiatzis D.G., McGranahan G.H. 1994 - *An improved method for acclimatizing tissue-cultured walnut plantlets using an antitranspirant*. Horticultural Science 29 (1): 4-2.
- Woeste K., Michler C. 2011- *Juglans*. In: "Forest trees Wild crop relatives: genomic and breeding resources". Chittaranjan K (ed), Springer, Berlin: 77-88.
- Yu Q., Mäntylä N., Salonen M. 2001 - *Rooting of Hybrid Clones of Populus tremula L. x P. tremuloides Michx. by Stem Cuttings Derived from Micropropagated Plants*. Scandinavian Journal of Forest Research 16 (3): 238-245.